

## Original Article

# On the effect of the injection of potassium phosphate *in vivo* inducing the precipitation of serum calcium with inorganic phosphate

Alcimar B Soares<sup>1</sup>, José G Ticianeli<sup>1</sup>, Letícia B M Soares<sup>2</sup>, George Amaro<sup>3</sup>

<sup>1</sup>Laboratory of Biomedical Engineering, Federal University of Uberlândia, Uberlândia, MG, Brazil; <sup>2</sup>Nephrology Service, General Hospital, Federal University of Uberlândia, Uberlândia, MG, Brazil; <sup>3</sup>Brazilian Agricultural Research Company, Boa Vista, RR, Brazil

Received October 25, 2013; Accepted November 30, 2013; Epub December 15, 2013; Published December 30, 2013

**Abstract:** High concentrations of inorganic phosphate (Pi) resulted from the hydrolysis of ATP is strongly associated to the weakness of the contractile mechanism of muscles due to its attractiveness to calcium. The majority of the experiments to study such effect are conducted *in vitro*. This work investigates the effects of different concentrations of Pi, induced by the injection of potassium phosphate in live animals, in the precipitation with serum calcium and the generation of calcium phosphate composites. The experiments were also designed to find out the ideal amount of potassium phosphate to induce an effective reaction. Potassium phosphate was injected in Wistar rats, randomly separated and distributed into seven groups. Group I was injected with 0.5 ml of saline solution (control) and groups II through VII were injected with 0.5, 1.5, 2.5, 5.0, 7.5 and 10.0 mg/kg of potassium phosphate, respectively. Blood collected from the inferior vena cava was submitted to biochemical analyses to measure the concentrations of calcium, Pi, urea and creatinine. The results showed that Pi, induced by the injection of potassium phosphate in live animals, causes precipitation with serum calcium, with statistically significant differences between the control and the treatment groups for doses up to 5.0 mg/kg. No statistically significant differences were found between the different doses and the concentration of urea and creatinine in the plasma. We conclude that potassium phosphate can be used to induce serum calcium precipitation *in-vivo*, with minor effects on other physiological variables, and the ideal dose to do so is 5.0 mg/kg.

**Keywords:** In vivo, potassium phosphate, inorganic phosphate, serum calcium, muscle fatigue

## Introduction

The effect of extracellular calcium and inorganic phosphate (Pi) on various aspects of the cellular mechanism has been the object of study of many research groups over the past decades [1-7]. Special attention has been devoted to the role of those substances in the underlying process of skeletal muscle contraction that begins with the release of a series of action potentials, promoting cell membrane depolarization and the release of calcium ions from the sarcoplasmic reticulum terminal [8-10]. Since the hydrolysis of adenosine triphosphate (ATP) is also required to provide energy and maintain the contraction, it is expected that the amount of ATP and its hydrolysis must be directly proportional to the demands of each contraction. Depending on the intensity of the contraction,

the hydrolysis of ATP can increase the concentration of cytoplasmic Pi up to 30 mM [11-13]. Although other biochemicals, including phosphocreatine (PCr), creatine, glucose 6-P and lactate also play an important role in the regulation of muscle contraction and power generation, high concentration of cytoplasmic Pi has been reported as one of the most important element contributing to fatigue and reduction of power [5, 11, 13, 14]. This can be explained by the attractiveness of Pi to calcium ions (Ca<sup>2+</sup>), released by the sarcoplasmic reticulum (SR), leading to the formation of an insoluble substance called calcium phosphate (CaPi) and reducing the quantities of Pi and Ca<sup>2+</sup> required to sustain the contraction [3].

Fryer *et al* were the first to suggest that Pi could enter the sarcoplasmic reticulum and precipi-

## Potassium phosphate *in vivo* induces precipitation of Pi with serum calcium

tate  $\text{Ca}^{2+}$ , forming CaPi and rapidly reducing the amount of calcium ions released, contributing to muscle fatigue [1]. Other works can also be found in the literature describing the effects of Pi reacting with  $\text{Ca}^{2+}$  released by the sarcoplasmic reticulum (SR) [4, 15-18]. However, since the majority of those experiments were carried out *in vitro*, it is possible that we are missing important metabolic reactions and substances, that could not be found or replicated *in vitro* but would play a significant role in the process of muscle contraction. Therefore, one can infer that the results of those surveys, while worthy of attention, are somehow limited to specific areas of interest. In fact, Fryer *et al* conclude that the omission of some possible *in vivo* reactions did not allow them to clearly establish if the decreased muscle strength was only due to precipitation of CaPi within the sarcoplasmic reticulum, or due to the loss of  $\text{Ca}^{2+}$  by the SR [1]. For example, cytoplasmic Pi can increase the probability of opening SR's ryanodine receptors (RyRs) [12, 19, 20]. RyRs are vital for muscle contraction since they are responsible for releasing  $\text{Ca}^{2+}$  into the cytosol [7, 21]. Hence, if cytoplasmic Pi can change the probability of opening RyRs channels, it can interfere with the whole process of muscle contraction. Also, cytoplasmic Pi may enter the SR and bind to  $\text{Ca}^{2+}$ , forming CaPi, further reducing the amount of releasable  $\text{Ca}^{2+}$  [6]. Nevertheless, it is possible that muscle strength is not affected by a significant reduction in  $\text{Ca}^{2+}$  available in the SR, since its endogenous contents of calcium (~1.1 mmol/l) may be much greater than necessary to fully activate the contractile mechanism [2, 9]. However, this is yet to be demonstrated.

In this article, we propose a method that can potentially be used to verify some of those important reactions *in vivo*. We investigate whether inorganic phosphate, induced by the injection of potassium phosphate ( $\text{K}_3\text{PO}_4$ ) in live animals, causes serum calcium precipitation, generating the insoluble calcium phosphate and, if so, what is the ideal amount to do it effectively. If the method proves to be feasible, without prejudice to the animal's health, it can contribute to the opening of new frontiers to the study of muscle fatigue in animals or humans.

### Materials and methods

#### Animals

Male rats of the Wistar strain, with an average age of 90 to 120 days, weighing between 50

and 150 gr, maintained on food and water *ad libitum*, housed in cages of white polyethylene (42x24x17 cm), with a maximum of five animals per cage, with daily hygiene, under controlled lighting conditions and light/dark cycle of 12/12 hours. The animals were acclimatized in the laboratory for at least 1 h. All experiments were conducted between 11:00 h and 17:00 h in order to minimize possible circadian variations [22].

Very old and very young animals or those presenting any disease that could affect the energetic or contractile capacity were excluded.

#### Groups

The animals were randomly separated and distributed into seven groups of seven animals each: Group I (control – sterile saline solution), Group II ( $\text{K}_3\text{PO}_4$ -0.5 mg/kg), Group III ( $\text{K}_3\text{PO}_4$ -1.5 mg/kg), Group IV ( $\text{K}_3\text{PO}_4$ -2.5 mg/kg), Group V ( $\text{K}_3\text{PO}_4$ -5.0 mg/kg), Group VI ( $\text{K}_3\text{PO}_4$ -7.5 mg/kg) and Group VII ( $\text{K}_3\text{PO}_4$ -10.0 mg/kg). As shown, the difference between the groups is the amount of potassium phosphate administered via rapid injection (for Groups II, III, IV, V, VI and VII), except for Group I, used as control and injected with sterile saline solution. In order to rule out the effect of animal handling or features of the vehicle used for drug administration, and to establish a base line for the analyses, the control group was injected with 0.5 ml of sterile saline solution (NaCl 0.9%).

It is important to emphasize that potassium phosphate is a substance widely used to control hypophosphatemia [23-27]. The amounts used in the experiments were based on the specifications given by Gaasbeek and Meinders [28], that used infusions of potassium phosphate at doses of 2.5 mg/kg.

After inoculation, the animals were kept in their cages for 10 minutes prior to the surgical procedure for blood sampling, with no restriction to movement. A previous pilot experiment performed by the authors showed that the concentrations of serum calcium, Pi, urea and creatinine were stable 10 minutes after the injection of different doses of  $\text{K}_3\text{PO}_4$ .

#### Surgical procedure

The animals were weighed and anesthetized with an intraperitoneal injection [29] of Sodium Pentothal 70 mg/kg (Anental®). They were considered anesthetized when there was no

## Potassium phosphate *in vivo* induces precipitation of Pi with serum calcium

response to mechanical stimuli and no movement was observed after a painful stimulus - gripping the tail. The animals were placed on a metal tray in the supine position, with pelvic and thoracic limbs restrained by adhesive tape. A median longitudinal incision, of approximately 8 cm, was made in the abdominal region with fine scissors, with exposure and lateral spacing of the bowel to the right, allowing the visualization of the inferior vena cava and withdrawal of venous blood, which was collected and conditioned in tubes without coagulants and centrifuged to obtain the serum (Bio Eng® Serum Centrifuge, Model BE-5000 no 1616). The serum was separated from the cellular portion of the blood no later than twenty minutes after collection, avoiding ion exchanges between the two due to the increased permeability of erythrocytes to calcium [30] and possible release of hematic phosphate [31], which could affect the results of the biochemical analysis. Hemolyzed blood samples were excluded from the biochemical analysis.

### Biochemical analysis

The concentrations of calcium, inorganic phosphate, urea and creatinine in the blood were measured threefold, and the biochemical values were used to represent the serum amounts of the substances. Blood samples were collected with Semi-automatic pipette (Digipet® 1000 µl) and analyzed by the following methods:

**Calcium:** *Cálcio-K007 (Bioclin®)* - End point colorimetric test.

This is a colorimetric test for *in vitro* diagnostic, based on the measured of the intensity of color produced by the compound formed by o-cresolphthalein complexone and calcium in alkaline pH. The concentration of calcium can be calculated by measuring the absorbance of the sample [32]. More details about this technique can be found in [30].

**Inorganic phosphate:** *Fósforo-K020 (Bioclin®)* - End point colorimetric test.

This test is based on the reaction of Pi with ammonium molybdate, forming ammonium phosphomolybdate, which is subsequently reduced to molybdenum-blue - a blue substance, which the intensity of color is proportional to the concentration of Pi [32]. By measuring the absorbance of the samples it is

possible to infer the concentration of Pi. A detailed description of the method is given by [31].

**Creatinine:** *Creatinina-K0106 (Bioclin®)* - End point colorimetric test (Jaffe modified).

It is a colorimetric test for *in vitro* diagnostic, based on the reaction of creatinine with picric acid, forming a reddish-yellow complex. In this pH occurs the maximum formation of the picric-creatinine color complex. By adding the acid reagent, pH is reduced and the color due to creatinine is undone, remaining the chromogens color [33]. The concentration of creatinine can be obtained by the difference between the absorbances at different pHs. A full description of the method can be found in [34].

**Urea:** *Uréia Enzimática-K047 (Bioclin®)* - End point enzymatic colorimetric test (Berthelot modified).

This test is based on the principle that urea can be hydrolyzed to ammonium ions and CO<sub>2</sub> by urease ( $urea + 3H_2O \xrightarrow{urease} 2NH_4^+ + CO_2 + 2OH^-$ ). At alkaline pH and in the presence of salicylate and sodium hypochlorite, ammonia reacts and results in a compound of green color whose intensity is proportional to the concentration of urea in the sample [33]. Therefore, by measuring the absorbance of the samples it is possible to infer the concentration of urea. A full description of method the can be found in [35].

The study was approved by the Ethics Committee for Animal Experimentation of the Federal University of Minas Gerais, Brazil, according to Protocol #020.

### Results

Dose-response curves were calculated for different doses of potassium phosphate versus the concentration of calcium, inorganic phosphate, urea and creatinine found in the plasma. The analyses of the dose-response curves were based on a graphical observation of the physiological responses and trend analysis. The tests of hypotheses were based on the average concentration of the substances in the blood, followed by a further analysis of trends. **Table 1** shows the Linear (1-2) and Nonlinear (3-5) models tested for trend analyses.

## Potassium phosphate *in vivo* induces precipitation of Pi with serum calcium

**Table 1.** Linear and nonlinear regression models tested when searching for best-fit method for the dose-response curves

1	Linear	$y_i = y_0 + ax$
2	Quadratic	$y_i = y_0 + ax + bx^2$
3	Negative exponential	$y_i = ae^{-bx}$
4	Negative exp. of three parameters	$y_i = y_0 + ae^{-bx}$
5	Sigmoidal of four parameters	$y_i = y_0 + \frac{a}{1 + e^{- \frac{X-X_0}{b} }}$

A model should be selected by taking into consideration its fitting to the data and also to the objectives of the study. In other words, prior to any consideration, a model would be selected according to: a) the biological logics - the objective, the nature, the scope and extension of the study, and the expected results; b) the significance of the adjustment of the model to the data; and c) the coefficient of determination ( $R^2$ ). If more than one model would apply - expressing a functional relationship between the studied variables and allowing an estimate of the parameters, capable of defining such relationship - the coefficient of determination and the biological logics expressed by them were also considered in order to select the most suitable model. In so doing, the chosen model would not only be a good statistical model, but also would generate the best representation of the biological aspects involved [36, 37].

Also, prior to the parametric analysis, each variable was tested for normality, independence of errors and homoscedasticity. The following procedure was used: (1) Kolmogorov-Smirnov normality test [38]; (2) Hartley's Fmax homoscedasticity test [39]; (3) Regression analyses with all models listed in **Table 1**; (4) Visual analysis of trends - adjustment of the model to the data, by means of a graph showing the mean values and errors.

After those tests and the preliminary graphic evaluation were concluded, it was found that all data could be represented by a normal distribution function (Calcium: K-S=0.12569,  $p < 0.20$ ; Inorganic phosphate: K-S=0.167401,  $p < 0.15$ ; Urea: K-S=0.11963,  $p < 0.20$ ; Creatinine: K-S=0.16510,  $p < 0.15$ ) and showed homogeneity of variance (Calcium: Fmax=2.9220,  $p < 0.1382$ ; Inorganic phosphate: Fmax=5.87,  $p <$

0.10; Urea: Fmax=2.2987,  $p < 0.1803$ ; Creatinine: Fmax=4.25,  $p < 0.10$ ).

### *Serum calcium concentration*

**Table 2** shows the estimated values for each parameter of the models, adjusted to the dose-response curve of serum calcium, according to the criteria described earlier. Note that all models show a good adjustment to the data ( $p < 0.05$ ). However, when

we compare the variation of the data obtained from the models and the total span of the real data, expressed by  $R^2$ , it was found that three of models showed a better adjustment (quadratic, sigmoidal and negative exponential). Nevertheless, the biological logic (the main criterion of choice) is based on the assumption that there may be a stabilization of calcium precipitation with inorganic phosphate. When testing such hypothesis, it was found that the sigmoidal regression is the best model to support it.

The graph in **Figure 1** represents the average of serum calcium for different doses of potassium phosphate. A reduction in the concentration of serum calcium can be observed as the doses of potassium phosphate increases up to 5.0 mg/kg, with high statistically significant differences. Above that value, there is a tendency towards stabilization of the response, with no statistically significant differences, as shown in **Table 3**. The continuous line shows the tendency (trend) estimated by the sigmoidal model.


**Table 3** shows the comparison between the mean values of serum calcium concentration for different doses of potassium phosphate. The values marked with '\*' are those that, according to the test, do not differ significantly, with a confidence level of 95%. Note that the control group (0.0 mg/kg of potassium phosphate) is significantly different from the others. The same can be seen for doses of 0.5, 1.5 and 2.5 mg/kg. Doses of 5.0, 7.5 and 10.0 mg/kg are statistically similar to each other, meaning that the concentration of serum calcium must have reached a stable level at 5.0 mg/kg of injected potassium phosphate. This effect can also be observed in **Figure 1**.

A similar strategy was used to evaluate the correlation between the doses of potassium phosphate and the concentration of inorganic phos-

## Potassium phosphate *in vivo* induces precipitation of Pi with serum calcium

**Table 2.** Estimated values for the parameters of the mathematical models, calculated for the serum calcium x potassium phosphate dose-response

Regression model	$Y_0$	a	b	$x_0$	$R^2$	$R^2_{aj}$	p
Linear	7.9430	-0.3180			0.7579	0.7095	0.0108
Quadratic	8.6239	-0.9604	0.0662		0.9841	0.9761	0.0003
Negative exponential		8.1247	0.0539		0.8090	0.7708	0.0058
Neg. exp. of three parameters	5.2387	3.5702	0.4046		0.9629	0.9444	0.0014
Sigmoidal of four parameters	5.3532	4.0662	-1.1411	1.5231	0.9744	0.9489	0.0069



**Figure 1.** Relationship between doses of potassium phosphate (mg/kg) and serum calcium (mg/dl) and the trend line (four-parameter sigmoidal model).

**Table 3.** Results of the Tukey test comparing the mean serum calcium (mg/dl) with different doses of potassium phosphate (mg/kg)

Potassium phosphate (mg/kg)	Serum calcium (mg/dl)
0.0	8.73
0.5	7.99
1.5	7.36
2.5	6.77
5.0	5.22*
7.5	5.37*
10.0	5.57*

Values marked with \*\*\* do not differ significantly, with confidence level of 95%.

phate, urea and creatinine in the plasma. The final results as shown next.

### Serum inorganic phosphate concentration

**Figure 2** shows the average concentration of inorganic phosphate (Pi) found in the blood for different doses of potassium phosphate. An increase in the concentration of Pi can be observed as the doses of potassium phosphate increases up to 2.5 mg/kg, with high statistically significant differences. Above that value, there is a tendency towards stabilization of the response, with no statistically significant differences, as shown in **Table 4**. The continuous line shows the tendency (trend) estimated by the sigmoidal model.


### Serum urea concentration

Although the data from urea analyses show a normal distribution and homogeneity of variance, none of the tested regression models showed statistical significance to represent the changes of urea as a function of the dosage of inorganic phosphate. Also, no statistically significant differences were found between different doses of potassium phosphate, with a confidence level of 95%. **Figure 3** shows the average of serum urea for different doses of potassium phosphate.

### Serum creatinine concentration

In the same way as for urea, although the creatinine analyses showed a normal distribution and homogeneity of variance, none of the tested regression models showed statistical signifi-

## Potassium phosphate *in vivo* induces precipitation of Pi with serum calcium



**Figure 2.** Relationship between doses of potassium phosphate (mg/kg) and serum Pi (mg/dl) and the trend line (four-parameter sigmoidal model).

**Table 4.** Results of the Tukey test comparing the mean serum Pi (mg/dl) for different doses of potassium phosphate (mg/kg)

Potassium phosphate (mg/ kg)	Serum Pi (mg/dl)
0.0	4.72
0.5	5.10
1.5	6.10
2.5	6.82*
5.0	7.06*
7.5	6.49*
10.0	6.66*

Values marked with \*\* do not differ significantly, with confidence level of 95%.

cance to represent the changes in serum concentration as a function of the dosage of inorganic phosphate. No statistically significant differences were found between different doses of potassium phosphate, with a confidence level of 95%. **Figure 4** shows the average of serum creatinine for different doses of potassium phosphate.

### Discussion

In this paper we have described a method for *in vivo* precipitation of calcium phosphate by


means of potassium phosphate inoculation. Unlike the *in vitro* approach, predominant for this kind of experiment, the method allowed us to avoid the omission of substances that could play important roles during *in vivo* experiments, as described earlier. Also, the injection of potassium phosphate can be done with minor effects on other physiological variables that could affect the desired results, allowing a new spectrum of researches to better understand the processes of muscle contraction and muscle fatigue.

The experimental setup consisted on the intraperitoneal injection of different doses of potassium phosphate and biochemical


analyses of blood collected from the inferior vena cava of the animals. Although we did not performed *in vitro* experiments, it is assumed that the precipitation of calcium phosphate occurred in a similar manner to the reactions reported by other *in vitro* experiments described in the literature [12, 16, 17].

The results show that injection of potassium phosphate raises the concentration of inorganic phosphate in the blood for doses up to 2.5 mg/kg - higher doses have no significant impact in the concentration of serum Pi. The amount of serum calcium was substantially reduced after the injection of potassium phosphate. Such reduction also varied according to the dosage of potassium phosphate, with no statistically significant changes for doses above 5.0 mg/kg. We assume that the dose-response has reached a certain metabolic equilibrium at 5.0 mg/kg. Confirming this assertion, the tests of trend analysis showed that the concentrations of serum calcium in response to inoculation of doses of 5.0, 7.5 and 10.0 mg/kg of potassium phosphate were basically the same (**Figure 1**, **Table 3**). When comparing the responses of calcium concentration between the control group (0.0 mg/kg of potassium phosphate) and the

## Potassium phosphate *in vivo* induces precipitation of Pi with serum calcium



**Figure 3.** Relationship between doses of potassium phosphate (mg/kg) and serum urea (mg/dl).



**Figure 4.** Relationship between doses of potassium phosphate (mg/kg) and serum creatinine (mg/dl).

Another important result gathered with this study was that the injection of potassium phosphate, at doses up to 10 mg/kg, did not have any significant impact in important metabolic functions during the experiments. It is known that changes in urea and creatinine may indicate abnormal functioning of the urinary system and protein metabolism [40]. Although there is some debate on the efficiency of those markers as indicators of premature renal damage, it is a consensus that high concentration of those substances in the plasma may indicate a variety of renal dysfunctions and damage [40, 41]. One might rightfully wonder whether high levels of urea could also be the result of the degradation of exogenous proteins from a feeding process. However, that was not the case of this study, since the animals were kept fasting prior to the experiments. High levels of creatinine could also be the result of intense physical activity or muscular lesions [42, 43], but again, to avoid such effect, the animals were not stressed prior to the experiments. Therefore, the absence of changes in the concentration of serum urea and creatinine (**Figures 3 and 4**) indicates that the injection of potassium phosphate did not have major effects on physiological variables that could be traced by urea or creatinine, for the duration of the experiments.

experimental Group V (5.0 mg/kg - promoter of the maximum effect, as shown in **Table 3**), a highly significant difference was observed.

By proving a method to induce the precipitation of calcium phosphate in live animals and defining and ideal dose to do so (5.0 mg/kg in rats),

# Potassium phosphate *in vivo* induces precipitation of Pi with serum calcium

we hope to provide a novel research tool for studying the role of inorganic phosphate and calcium concentration in various physiological processes, in particular, those related to skeletal muscle fatigue.

## Acknowledgements

The authors would like to thank Professor MSc. Sabrina Degaspore for the important contribution to the experiments; Professor Kelly Becker Cristiane Surian for performing the biochemical analysis and the academics Wallace Bruno Ferreira Garcia and Sebastian Salazar Jansen Filho for their help during the experiments; to FAPEMIG-MG-Brasil, CNPq-Brasil and CAPES-Brasil for the financial support of this research.

## Disclosure of conflict of interest

None.

**Address correspondence to:** Dr. Alcimar B Soares, Laboratory of Biomedical Engineering, Federal University of Uberlândia, Uberlândia, MG, Brazil. E-mail: alcimar.soares@gmail.com

## References

- [1] Fryer MW, Owen VJ, Lamb GD and Stephenson DG. Effects of creatine phosphate and Pi on Ca<sup>2+</sup> movements and tension development in rat skinned skeletal muscle fibres. *J Physiol* 1995; 482: 123-140.
- [2] Owen VJ, Lamb GD, Stephenson DG and Fryer MW. Relationship between depolarization-induced force responses and Ca<sup>2+</sup> content in skeletal muscle fibres of rat and toad. *J Physiol* 1997; 498: 571-586.
- [3] Westerblad H, Bruton JD, Allen DG and Lännergren J. Functional significance of Ca<sup>2+</sup> in long-lasting fatigue of skeletal muscle. *Eur J Appl Physiol* 2000; 83: 166-174.
- [4] Duke A and Steele D. Mechanisms of reduced SR Ca<sup>2+</sup> release induced by inorganic phosphate in rat skeletal muscle fibers. *Am J Physiol Cell Physiol* 2001; 281: C418-C429.
- [5] Tesi C, Colomo F, Piroddi N and Poggesi C. Characterization of the cross-bridge force-generating step using inorganic phosphate and BDM in myofibrils from rabbit skeletal muscles. *J Physiol* 2002; 541: 187-199.
- [6] Dutka TL, Cole L and Lamb GD. Calcium phosphate precipitation in the sarcoplasmic reticulum reduces action potential-mediated Ca<sup>2+</sup> release in mammalian skeletal muscle. *Am J Physiol Cell Physiol* 2005; 289: C1502-C1512.
- [7] Amador FJ, Stathopoulos PB, Enomoto M and Ikura M. Ryanodine receptor calcium release channels: lessons from structure-function studies. *FEBS J* 2013; 280: 5456-5470. doi: 10.1111/febs.12194.
- [8] Borejdo J, Szczesna-Corday D, Muthu P, Metticolla P, Luchowski R, Gryczynski Z and Gryczynski I. Single Molecule Detection Approach to Muscle Study: Kinetics of a Single Cross-Bridge During Contraction of Muscle. *Methods Mol Biol* 2012; 875: 311-334. doi: 10.1007/978-1-61779-806-1\_17.
- [9] Yanagida T and Ishii Y. Muscle contraction mechanism based on single molecule measurements. *J Muscle Res Cell Motil* 2012; 33: 395-402.
- [10] Berridge MJ, Lipp P and Bootman MD. The Versatility and Universality of Calcium Signaling. *Nat Rev Mol Cell Biol* 2000; 1: 1-21.
- [11] Yin Y, Guo Z, Chen X, Xing C and Fan Y. Studies on biomechanics of skeletal muscle based on the working mechanism of myosin motors: An overview. *Chin Sci Bull* 2012; 57: 4533-4544.
- [12] Balog EM, Fruen BR, Kane PK and Louis CF. Mechanisms of Pi regulation of the skeletal muscle SR Ca<sup>2+</sup> release channel. *Am J Physiol Cell Physiol* 2000; 278: C601-C611.
- [13] Debold EP, Dave H and Fitts RH. Fiber type and temperature dependence of inorganic phosphate: implications for fatigue. *Am J Physiol Cell Physiol* 2004; 287: C673-C681.
- [14] He Z, Chillingworth RK, Brune M, Corrie JET, Webb MR and Ferenczi MA. The efficiency of contraction in rabbit skeletal muscle fibres, determined from the rate of release of inorganic phosphate. *J Physiol* 2003; 517: 839-854.
- [15] Allen DG and Westerblad H. Role of phosphate and calcium stores in muscle fatigue. *J Physiol* 2001; 536: 657-665.
- [16] Amrute-Nayak M, Antognozzi M, Scholz T, Kojima H and Brenner B. Inorganic phosphate binds to the empty nucleotide binding pocket of conventional myosin II. *J Biol Chem* 2008; 283: 3773-3781.
- [17] Debold EP, Longyear TJ and Turner MA. The effects of phosphate and acidosis on regulated thin-filament velocity in an *in vitro* motility assay. *J Appl Physiol* 2012; 113: 1413-1422. doi: 10.1152/jappphysiol.00775.2012.
- [18] Duke A and Steele D. Interdependent effects of inorganic phosphate and creatine phosphate on sarcoplasmic reticulum Ca<sup>2+</sup> regulation in mechanically skinned rat skeletal muscle. *J Physiol* 2001; 531: 729-742.
- [19] Fruens BR, Mickelson JR, Shomer NH, Roghair TJ and Louis CF. Regulation of the Sarcoplasmic Reticulum Ryanodine Receptor by Inorganic Phosphate. *J Biol Chem* 1994; 269: 192-196.



## Potassium phosphate *in vivo* induces precipitation of Pi with serum calcium

- [20] Rannou F, Leschiera R, Giroux-Metges MA and Pennec JP. Effects of lactate on the voltage-gated sodium channels of rat skeletal muscle: modulating current opinion. *J Appl Physiol* 2012; 112: 1454-1465. doi:10.1152/jap-physiol.00944.2011.
- [21] Fill M and Copello JA. Ryanodine receptor calcium release channels. *Physiol Rev* 2002; 82: 893-922.
- [22] Bittman EL and Johnson CH. Animal Care Practices in Research on Biological Rhythms and Sleep. *J Biol Rhyth* 2013; 28: 303-304. doi: 10.1177/0748730413502313.
- [23] Basri MN, Janattul AJ, Azrina MR and Abdul Hadi M. Hypophosphatemia in the Intensive Care Unit: Incidence, Predictors and Management. *Med J Malaysia* 2012; 11: 31-36.
- [24] Liamis G, Milionis HJ and Elisaf M. Medication-induced hypophosphatemia: A review. *Q J Med* 2010; 103: 449-459. doi:10.1093/qjmed/hcq039.
- [25] Miller DW and Slovis CM. Hypophosphatemia in the Emergency Department Therapeutics. *Am J Emerg Med* 2000; 18: 457-461.
- [26] Sheldon GF. Treatment of Hypophosphatemia. *J Am Coll Surg* 2004; 199: 171-181.
- [27] Taylor BE, Huey WY, Buchman TG, Boyle WA and Coopersmith CM. Treatment of Hypophosphatemia using a protocol based on patient weight and serum phosphorus level in a surgical intensive care unit. *J Am Coll Surg* 2004; 198: 198-204.
- [28] Gaasbeek A and Meinders E. Hypophosphatemia: An update on its etiology and treatment. *AM J Med* 2005; 118: 1094-1101.
- [29] Blouin A, Molez S, Pham D, Ayach B, Dussault B, Escher E, Jeng A and Battistini B. A Novel Procedure for daily measurements of hemodynamical, hematological, and biochemical parameters in conscious unrestrained rats. *J Pharmacol Toxicol Meth* 2000; 44: 489-505.
- [30] Bioclin. Cálcio-K007 – Instruções de uso. Available from: [http://www.bioclin.com.br/sitebioclin/wordpress/wp-content/uploads/arquivos/instrucoes/instrucoes\\_calcio\\_colorimetrico.pdf](http://www.bioclin.com.br/sitebioclin/wordpress/wp-content/uploads/arquivos/instrucoes/instrucoes_calcio_colorimetrico.pdf). Accessed on: 01 August 2013. 2013.
- [31] Bioclin. Fósforo-K020 – Instruções de uso. Available from: [http://www.bioclin.com.br/sitebioclin/wordpress/wp-content/uploads/arquivos/instrucoes/instrucoes\\_fosforo.pdf](http://www.bioclin.com.br/sitebioclin/wordpress/wp-content/uploads/arquivos/instrucoes/instrucoes_fosforo.pdf). Accessed on: 01 August 2013. 2013.
- [32] Cançado JR, Lima AO, Soares JB, Grego JB and Galizi J. *Métodos de Laboratórios Aplicados a Clínica*. 8ª ed. Rio de Janeiro: Guanabara Koogan, 2001.
- [33] Henry JB. *Diagnósticos Clínicos e Tratamento por Métodos Laboratoriais*. 21ª ed. São Paulo: Editora Manole, 2013.
- [34] Bioclin. Creatinina-K016 – Teste colorimétrico de ponto final (Jaffé Modificado) ou cinética de tempo fixo. Available from: [http://www.bioclin.com.br/sitebioclin/wordpress/wp-content/uploads/arquivos/instrucoes/instrucoes\\_creatinina.pdf](http://www.bioclin.com.br/sitebioclin/wordpress/wp-content/uploads/arquivos/instrucoes/instrucoes_creatinina.pdf). Accessed on: 01 August 2013. 2013.
- [35] Bioclin. Uréia enzimática-K047 – Enzimático colorimétrico de ponto final (Berthelot Modificado) ou cinética de tempo fixo. Available from: [http://www.bioclin.com.br/sitebioclin/wordpress/wp-content/uploads/arquivos/instrucoes/instrucoes\\_ureia\\_enzimatica.pdf](http://www.bioclin.com.br/sitebioclin/wordpress/wp-content/uploads/arquivos/instrucoes/instrucoes_ureia_enzimatica.pdf). Accessed on: 01 August 2013. 2013.
- [36] Dell RB, Holleran S and Ramakrishnan R. Sample Size Determination. *ILAR J* 2002; 43: 207-213.
- [37] Bartlett JE, Kotrlik JW and Higgins C. Organizational research: Determining appropriate sample size for survey research. *Inf Technol Learn Perform J* 2001; 19: 43-50.
- [38] Zar JH. *Biostatistical analysis*. 5th edition. New Jersey: Prentice Hall, 2010.
- [39] Indrayan A. *Medical Biostatistics*. 3rd edition. New York: CRC Press, 2012.
- [40] Siew ED, Ware LB and Ikizler TA. Biological Markers of Acute Kidney Injury. *J Am Soc Nephrol* 2011; 22: 810-820. doi: 10.1681/ASN.2010080796.
- [41] Salgado JV, Neves FA, Bastos MG, França AK, Brito DJ, Santos EM and Salgado Filho N. Monitoring renal function: measured and estimated glomerular filtration rates - a review. *Braz J Med Biol Res* 2010; 43: 528-536.
- [42] Kopp J, Loos B, Spilker G and Horch RE. Correlation between serum creatinine kinase levels and extent of muscle damage in electrical burns. *Burns* 2004; 30: 680-683.
- [43] Zhu BL, Ishida K, Quan L, Taniguchi M, Oritani S, Li DR, Fujita MQ and Maeda H. Postmortem serum uric acid and creatinine levels in relation to the causes of death. *Forensic Sci Int* 2002; 125: 59-66.