



ESTs and putative line-specific (broiler and layer) SNPs identified in genes expressed in *Gallus gallus* pituitary and hypothalamus

Clarissa Sanches da Silva Cassoli¹, Erika Cirstina Jorge¹, Mateus Patricio¹, Helena Javiel Alves¹, Erik Amazonas de Almeida¹, Mônica Corrêa Ledur² and Luiz Lehmann Coutinho¹

¹Laboratório de Biotecnologia Animal, Departamento de Zootecnia,

Escola Superior de Agricultura "Luiz de Queiroz", Universidade de São Paulo, Piracicaba, SP, Brazil.

²EMBRAPA Suínos e Aves, Concórdia, SC, Brazil.

Abstract

Brazilian poultry industry has reached a high level of development in both meat and egg production as a result of constant technological modernization. Further improvements can be achieved through genomics, but before this can be accomplished, a better understanding of gene expression profiles and nucleotide polymorphisms is necessary. Since animal physiology is directly or indirectly controlled by the pituitary and hypothalamus, the aim of the present work was to identify and analyze genes expressed in these tissues in chicken lines with different growth potential. Two pituitary and hypothalamus cDNA libraries from 21 day broiler (TT) and layer (CC) chickens lines were constructed and allowed identification of 3,074 unique sequences and 77 single nucleotide polymorphisms (SNPs). The collection of expressed sequence tags (ESTs) and SNPs identified in this study represents an important resource for future studies aimed at identifying genes responsible for growth in chicken.

Key words: pituitary, hypothalamus, EST, SNP, chicken.

Received: February 1, 2007; Accepted: November 20, 2007.

Introduction

Over the last decades, the poultry industry has experienced a substantial increase in production efficiency. For example, there has been a threefold increase in egg production per chicken/year and a substantial decrease in the time necessary for broilers to reach 1.5 kg of live weight (Burt, 2002). Classic selection greatly contributed to this progress, since selected broiler lines grow 3 to 4 times more rapidly than their non-selected ancestor, red jungle fowl (Bulfield, 2004). Nevertheless, several phenotypic traits are difficult to improve through traditional breeding, such as those that are difficult or expensive to measure (carcass quality and composition, behavior and welfare) or have low heritability (reproduction and fitness) (Bulfield, 2004). In addition, unwanted characteristics can be indirectly selected with classic breeding (Burt, 2002). Modern genomic technologies can greatly impact selection of these difficult to target traits (Bulfield, 2004).

Important aspects of animal physiology are directly or indirectly controlled by the pituitary and hypothalamus,

but the genetic mechanisms controlling processes such as metabolism, somatic growth and reproduction in chickens still remain largely unknown (Cogburn *et al.*, 2003). Therefore, identification and study of genes expressed in the pituitary and hypothalamus can fill existing gaps in understanding the molecular pathways involved in several physiological processes, as well as provide tools for future animal breeding programs.

Among the methodologies available for gene identification, analysis of expressed sequence tags (ESTs) has proven to be very efficient. This methodology consists in partially sequencing the extremities of clones obtained from cDNA libraries and establishing groups of specifically expressed genes, as well as their transcription levels in determined tissue or cell types (Adams *et al.*, 1991). This approach allows comparisons to be made between different tissues or species, and the identification of polymorphisms in intragenic regions (Adams *et al.*, 1991; Hatey *et al.*, 1998). Single nucleotide polymorphisms (SNPs) have emerged as a principal DNA marker class which greatly helps in developing high-density genetic maps for use in QTL identification through linkage disequilibrium analysis (Smith *et al.*, 2002).

In previous studies, ESTs were generated from skeletal muscle precursor tissues (somites and neural tube),

Send correspondence to Luiz Lehmann Coutinho. Laboratório de Biotecnologia Animal, Departamento de Zootecnia, Escola Superior de Agricultura "Luiz de Queiroz", Universidade de São Paulo, 13418-900 Piracicaba, SP, Brazil. E-mail: llc@usp.br.

limbs and whole embryos (Jorge *et al.*, 2004), and from young fowl pectoral musculature (unpublished results). The present work was developed with the scope of cataloguing genes expressed in chicken hypothalamus and pituitary gland and identifying distinct features possibly associated with growth.

Material and Methods

A total of about 120 eggs from broiler (TT) and layer (CC) lines supplied by Embrapa Swine and Poultry National Research Center were incubated at 37 °C in a humidity-controlled atmosphere. The TT line is a male line obtained from a cross of *Cornish*, *Hampshire* and *Plymouth Rock* breeds. This line has been selected for meat production since 1985. In the first stages of the breeding program the focus was upon weight gain and carcass traits; however, since 1992, males of this lineage have also been selected for feed conversion rate. CC line is a female line of White Leghorn selected initially (1989) for egg production and quality. After hatching, chicks were kept in a commercial broiler house at the Animal Science Department, ESALQ-USP. Chicks were initially given commercial broiler feed and water *ad libitum*. They were exposed to room temperature and continuous luminosity until 21 days of age. At this age, the pituitary and hypothalamus were extracted surgically and stored in liquid nitrogen. Total RNA was extracted from each line separately according to the protocol described by Chomczynski and Sacchi (1987), followed by poly(A)+RNA isolation using the Oligotex kit (GE HealthCare).

cDNA libraries were constructed from 1-2 µg poly(A)+ RNA using the SuperScript Plasmid System kit (Invitrogen), according to manufacturer's protocol. Fractions containing cDNA larger than 500 bp were ligated into the Sall-NotI site of pSPORT1 vector (Invitrogen). Clones were sequenced from the 5' ends using the Big Dye Terminator Cycle Sequencing Ready Reaction kit (Applied Biosystems) in conjunction with primer T7 (5'-TAATACGACTCATATAGGG-3').

Sequences were analyzed for nucleotide quality with PHRED software (Ewing *et al.*, 1998). Sequences considered valid (minimum of 150 bp with quality > 20) were grouped with the CAP3 software (Huang and Madan, 1999). Unique sequences (contigs + singletons) were compared to the non-redundant (nr) GenBank database (www.ncbi.nlm.nih.gov) using the Blastx algorithm (Altschul *et al.*, 1990).

Comparative analysis of expression patterns was performed between the two lines and all other libraries constructed in the laboratory, according to the relative frequency of the ESTs. SNPs were identified analyzing nucleotide discrepancies between ESTs from the two lines. Only those SNPs that appeared at least twice in the same line/posi-

sition and showed PHRED quality equal or higher than 20 were considered as hypothetic line-specific SNPs.

Results and Discussion

Sequence analysis

A total of 5,017 ESTs were obtained from the 5' end of the inserts cloned in the two cDNA libraries. By PHRED analysis, 2,133 reads from broiler line (TT) and 2,153 of the layer line (CC) were considered valid. Sequences referring to the CC line library were deposited in dbEST division of GenBank with accession numbers ranging from CO419474 to CO421626, and those referring to the TT line library received numbers ranging from CO421627 to CO423759. After clustering and assembly (CAP3 software, Huang and Madan, 1999), the TT line library presented 1,643 unique sequences (contigs + singletons). Of these, 1,477 were singletons and 656 were grouped into 166 contigs, whose sequence number varied from 2 to 59. The CC line library represented 1,649 unique sequences, with 1,475 singletons and 678 sequences grouped into 174 contigs, with 2 to 70 sequences per contig. This clustering indicated novelty rates of 77% for the TT library and 76.6% for the CC line library. Both library sequences were also clustered together, revealing 3,074 unique sequences and a novelty index of 71.1%.

Expression profile ("digital Northern")

Relative frequency of ESTs was compared using all *G. gallus* sequences obtained in the laboratory (a total of 13,521 ESTs), in a strategy known as "digital Northern" (Audic and Claverie, 1997). The clustering and assembly of all 13,521 ESTs resulted in 680 contigs formed by grouping TT and CC ESTs. Sequences present in 472 (69.4%) of these contigs were also identified in libraries constructed from somites, limbs (Jorge *et al.*, 2004) and breast muscle, suggesting that these ESTs are coordinately expressed in all the different tissues studied. Sequences present in 94 contigs (13.8%) were identified as unique to the pituitary and hypothalamus libraries in both TT and CC lines, and were called library-specific contigs. The 114 remaining contigs (16.8%) were library and line-specific, since they were only encountered in the pituitary and hypothalamus libraries of the lines studied.

Among the ESTs identified as library and library/line specific, sequences coding for proteins known to participate in molecular pathways for growth and reproduction were identified. Some examples of genes represented preferentially in the CC line were Ca²⁺/Calmodulin-dependent protein kinases proteins (CAMK1 and CAMK2), G protein-coupled receptors (GPRs), cAMP phosphodiesterase (LOC771318, LOC425199), phosphoinositide 3-kinase (PIK3), and N-myc downstream-regulated gene 1 protein (NDR1).

SNPs identification

Following clustering and assembly of the sequences from the pituitary and hypothalamus libraries with CAP3 software, ESTs grouped in the same contig were used to search for single nucleotide polymorphisms. Only those SNPs that appeared at least twice in the same line/position and showed PHRED quality equal or higher than 20 were considered as hypothetical line-specific SNPs.

Of the 389 contigs constituted by sequences from the pituitary and hypothalamus libraries, 28 (7.2%) presented 77 line-specific SNPs, corresponding to 52 TT-specific and 25 CC-specific SNPs (Table 1). Most SNPs found in ESTs were related to the mitochondrial genome, to structural proteins, neuronal constituents, ribosomal proteins and iron binding proteins. SNPs were also observed in hypothetical proteins (proteins still lacking a defined biological function), calcium binding proteins, lipid metabolism related

proteins and ESTs lacking similarity to any sequence in the database consulted.

PHRED quality of the SNP nucleotides varied from 20 to 68, with an average value of 50.1. Among the 28 contigs, 14 presented only one SNP, six presented two SNPs, three showed three SNPs, one contig presented five SNPs, one had seven SNPs, two contigs showed eight SNPs and one presented 14 SNPs. These 28 contigs were constituted by four to 112 ESTs each, presenting an average number of 21.2 ESTs per contig and a mean length of 1,441 bp. Minimum length was 746 bp and maximum length was 3,006 bp. Contig SNP density varied from 0.5 to 9.1 SNPs per kb, with an average density of 1.9 SNP/kb. This result is within the estimated polymorphism range observed by Smith *et al.* (2002). TT-specific SNPs showed greater density than the CC-specific ones. Average density values were 1.3 and 0.6, respectively. This two times higher density value for TT-specific SNPs is interesting since the total

Table 1 - Relation of putative line specific SNPs found in 28 contigs.

Accession number /unigene cluster	Blast hit	ESTs number of the cluster/cluster size	Consensus	Mutation (line - number of occurrence)	Position (PHRED quality)
CO419502 Gga.2913	MBP myelin basic protein	112 (52 CC + 60 TT) 2,739 bp	A	G (TT - 2)	61 (q:30,30)
			A	C (TT - 2)	67 (q: 22, 24)
			T	A (CC - 2)	72 (q: 43, 53)
			C	G (CC - 2)	73 (q: 43, 53)
			A	T (TT - 2)	80 (q: 33, 57)
			G	C (TT - 2)	99 (q: 53, 47)
			G	A (TT - 2)	460 (q: 30, 20)
			A	G (CC - 5)	942 (q: 57, 62, 57, 62, 39)
					1008 (q: 62, 52, 20, 68)
			C	T (TT - 4)	1494 (q: 68, 47, 62, 28, 50)
					1541 (q: 20, 53, 68)
			T	C (TT - 5)	1680 (q: 68, 68, 68, 21)
					1818 (q: 52, 39)
			T	C (CC - 3)	1872 (q: 20, 22)
CO419476	COX1 cytochrome c oxidase sub-unit I (mitochondrial protein)	90 (49 CC + 41 TT) 1,703 bp	A	T (CC - 2)	37 (q: 50, 53)
			A	T (CC - 2)	672 (q: 36, 22)
CO419511	Hypothetical protein (mitochondrial protein)	78 (68 CC + 10 TT) 1,862 bp	T	G (CC - 4)	61 (q: 43, 30, 37, 35)
					62 (q: 22, 43, 37, 42, 42)
			T	C (CC - 5)	63 (q: 27, 53, 44, 42, 43)
					64 (q: 54, 53, 53, 47, 50)
			A	G (CC - 5)	67 (q: 57, 57, 53, 57)
			A	T (CC - 5)	
CO419506 Gga.13261	similar to Myelin proteolipid protein (PLP) (Lipophilin)	35 (17 CC + 18 TT) 1,356 bp	T	G (CC - 4)	
			A	C (TT - 2)	106 (q: 47, 68)
			G	A (TT - 2)	107 (q: 36, 68)
			G	C (TT - 2)	110 (q: 53, 68)
			A	T (TT - 2)	112 (q:57, 57)
			A	C (TT - 2)	114 (q: 59,57)
			G	C (CC - 3)	128 (q: 36, 33, 33)
					132 (q: 50, 50, 42)
			A	G (CC - 3)	137 (q:47, 47)
			G	C (CC - 2)	

Table 1 (cont.)

Accession number /unigene cluster	Blast hit	ESTs number of the cluster/cluster size	Consensus	Mutation (line - number of occurrence)	Position (PHRED quality)
CO419546	COX3 cytochrome c oxidase subunit III (mitochondrial protein)	32 (14 CC + 18 TT) 805 pb	A C	C (TT - 3) G (TT - 2)	17 (q: 47,47,42) 19 (q: 53, 47)
CO421660 Gga.13261	similar to Myelin proteolipid protein (PLP) (Lipophilin)	27 (14 CC + 13 TT) 2,059 bp	A C	G (TT - 3) G (TT - 2)	1005 (q: 41, 68, 26) 1877 (q: 40, 46)
CO421668	CYTB cytochrome b (mitochondrial protein)	20 (8 CC + 12 TT) 1,185 bp	A	T (CC - 2)	729 (q:22, 22)
CO419634 Gga.6482	Heat shock protein HSP 90-alpha	20 (9 CC + 11 TT) 3,006 bp	G C	A (TT - 2) T (TT - 2)	119 (q: 47, 68) 365 (q: 62, 68)
CO419555	Hypothetical protein	19 (10 CC + 9 TT) 1,814 bp	T	C (CC - 2)	1158 (q: 54, 68)
CO421863 Gga.35012	Tubulin, alpha 1c	17 (6 CC + 11 TT) 1,343 bp	C	T (TT - 2)	238 (q: 32, 62)
CO421647	Hypothetical protein	15 (2 CC + 13 TT) 1,418 bp	G G	A (TT - 2) A (TT - 3)	125 (q:68, 68) 434 (q: 68, 68, 62) 466 (2: 32, 47) 767 (q: 43, 39) 780 (q: 32, 27) 885 (q: 59, 30) 907 (q: 57, 35)
CO422016 Gga.39142	UBC ubiquitin C	14 (9 CC + 5 TT) 880 bp	T C G T A G	C (CC - 3) T (TT - 2) A (TT - 2) C (TT - 2) C (TT - 2) A (CC - 3)	105 (q: 35, 68, 68) 174 (q: 68, 27) 249 (q: 68, 44) 264 (q: 57, 41) 354 (q: 50, 68) 447 (q: 39, 48, 50) 477 (q: 37, 35) 537 (q: 24, 46, 37)
CO422026	ND4L NADH dehydrogenase subunit 4L	11 (2 CC + 9 TT) 1,425 bp	T	C (CC - 2)	667 (2: 62, 51)
CO421775 Gga.34313	Apolipoprotein A-I	11 (4 CC + 7 TT) 1,302 bp	C	G (TT - 4)	459 (q: 42,48, 50, 52)
CO422318 Gga.1383	Enolase 1, (alpha)	10 (4 CC + 6 TT) 1,667 bp	C T C	T (TT - 4) C (TT - 2) T (TT - 4)	516 (q: 52, 59, 68, 41) 747 (q: 22, 21)
CO421930 Gga.2613	SPARC Secreted protein, acidic, cysteine-rich (osteonectin)	9 (7 CC + 2 TT) 1,719 bp	T C A	A (TT - 3) T (TT - 2) T (TT - 2) G (TT - 2)	283 (q: 68, 42) 289 (q: 68, 38, 26, 68) 340 (q: 68, 57, 68)
CO419533 Gga.26418	Calmodulin 2 (phosphorylase kinase, delta)	9 (5 CC + 4 TT) 1,560 bp	C	T (TT - 2)	989
CO421674 Gga.22905	TUBB2C tubulin, beta 2C	8 (3 CC + 5 TT) 746 bp	A	G (TT - 2)	47
CO421914 Gga.37995	similar to Neuron specific protein family member 2 (Protein p19) (HMP19)	7 (3 CC + 4 TT) 1,544 bp	T	C (TT - 2)	967 (q: 35, 41)
CO421925 Gga.35414	Neurofilament, light polypeptide 68kDa	7 (2 CC + 5 TT) 893 bp	C	T (TT - 2)	66 (q: 66, 66)

Table 1 (cont.)

Accession number /unigene cluster	Blast hit	ESTs number of the cluster/cluster size	Consensus	Mutation (line - number of occurrence)	Position (PHRED quality)
CO422566 Gga.4080	Ribosomal protein L7a	7 (3 CC + 4 TT) 883 bp	T	C (CC - 2)	282 (q: 62, 68)
CO419647 Gga.4827	Solute carrier family 25 (mitochondrial carrier; adenine nucleotide translocator), member 6	7 (3 CC + 4 TT) 2,741 bp	C	T (CC - 2)	459 (q: 68, 68)
CO422015 Gga.33759	Eukaryotic translation elongation factor 1 alpha 2	5 (1 CC + 4 TT) 866 bp	C	T (TT - 2)	381 (q: 57, 62)
CO422299		5 (2 CC + 3 TT) 873 bp	A G	C (TT - 2) A (CC - 2)	540 (q: 22, 50) 566 (q: 43, 54)
CO422068 Gga.35012	Tubulin, alpha 1c	5 (2 CC + 3 TT) 1,258 bp	C	T (TT - 2)	651 (q: 23, 33)
CO420128 Gga.1141	Malate dehydrogenase 1, NAD (soluble)	5 (3 CC + 2 TT) 1,428 bp	T G	C (TT - 2) A (TT - 2)	222 (q: 44, 41) 255 (q: 35, 52)
CO422266 Gga.21300	Similar to CD59 protein	4 (2 CC + 2 TT) 815 bp	A	C (TT - 2)	355 (q: 57, 57)
CO419495 Gga.12910	Angiotensinogen (serpin peptidase inhibitor, clade A, member 8)	4 (1 CC + 3 TT) 1,071 bp	G	C (TT - 2)	726 (q: 48, 37)

number of ESTs in the TT contigs was actually slightly lower than in CC (305 SNPs in ESTs from CC library versus 288 ESTs in the TT). This fact suggests that the TT line is more polymorphic, presenting higher allele numbers than the CC line.

These putative line-specific SNPs were classified according to the nucleotide substitution as transitions (purine → purine or pyrimidine → pyrimidine) or transversions (purine → pyrimidine → purine). Of the 77 identified SNPs, 50 (64.9%) were classified as transitions, with 36 (46.7%) being TT-specific and 14 (18.2%) CC-specific; 27 (35.1%) were classified as transversions, with 16 (20.8%) being TT-specific SNPs and 11 (14.3%) CC-specific. This finding gives a transition/transversion ratio of 1.85:1.

The EST collection obtained in this study allowed the identification of genes expressed in the pituitary and hypothalamus in two commercial chicken lines, thus providing an important resource for studies of growth physiology and animal breeding. In addition, we identified a series of line-specific SNPs. The SNPs identified in genes specifically expressed in this major control center of animal physiology hold great potential for selection studies, since SNPs are the most frequent form of genome variation and are currently considered a new generation of molecular markers.

References

- Adams MD, Kelley JM, Gocayne JD, Dubnick M, Polymeropoulos MH, Xiao H, Merril CR, Wu A, Olde B, Moreno RF, et al. (1991) Complementary DNA sequencing: Expressed sequence tags and human genome project. *Science* 252:1651-1656.
- Altschul SF, Gish W, Miller W, Myers EW and Lipman DJ. (1990) Basic local alignment search tool. *J Mol Biol* 215:403-410.
- Audic S and Claverie JM (1997) The significance of digital gene expression profile. *Genome Res* 7:986-995.
- Bulfield G (2004) Poultry breeding in the post-genomics era. *Br Poult Sci* 45:5-8.
- Burt DW (2002) Applications of biotechnology in the poultry industry. *Worlds Poult Sci J* 58:5-13.
- Chomczynski P and Sacchi N (1987) Single step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem* 162:156-159.
- Cogburn LA, Wang X, Carre W, Rejto L, Porter TE, Aggrey SE and Simon J (2003) Systems-wide chicken DNA microarrays, gene expression profiling, and discovery of functional genes. *Poult Sci* 82:939-951.
- Ewing B, Hillier L, Wendl MC and Green P (1998) Base-calling of automated sequencer traces using Phred. I. Accuracy assessment. *Genome Res* 8:175-185.
- Hatey F, Tosser-Klopp G, Cloucard-Martino C, Mulsant P and Gasser F (1998) Expressed sequence tags for genes: A review. *Genet Select Evol* 30:521-541.
- Huang X and Madan A (1999) Cap3: A DNA sequence assembly program. *Genome Res* 9:868-877.
- Jorge EC, Monteiro-Vitorello CB, Alves HJ, Silva CS, Balan RG, Patricio M and Coutinho LL (2004) EST analysis of mRNAs expressed during embryogenesis in *Gallus gallus*. *Int J Dev Biol* 48:333-337.
- Smith EJ, Shi L and Smith G (2002) Expressed sequence tags for the chicken genome from a normalized 10-day-old white leghorn whole-embryo cDNA library. 3. DNA sequence analysis of genetic variation in commercial chicken populations. *Genome* 45:261-267.

Assistant Editor: Klaus Hartfelder